


☐

I'm not robot


reCAPTCHA

Continue

How quickly should docusate sodium work

Stool softeners come as a capsule, tablet, liquid, and syrup to take by mouth. A stool softener usually is taken at bedtime. Follow the directions on the package or your prescription label carefully, and ask your doctor or pharmacist to explain any part you do not understand. Take stool softeners exactly as directed. Do not take more or less of it or take it more often than prescribed by your doctor.Swallow the docusate capsules whole; do not split, chew, or crush them.Take capsules and tablets with a full glass of water. The liquid comes with a specially marked dropper for measuring the dose. Ask your pharmacist to show you how to use it if you have difficulty. Mix the liquid (not the syrup) with 4 ounces (120 milliliters) of milk, fruit juice, or formula to mask its bitter taste.One to three days of regular use usually are needed for this medicine to take effect. Do not take stool softeners for more than 1 week unless your doctor directs you to. If sudden changes in bowel habits last longer than 2 weeks or if your stools are still hard after you have taken this medicine for 1 week, call your doctor. DOCUSATE SODIUM; SENNA (doc CUE sayt SOE dee um; SEN na) contains a stool softener and a laxative. It is used to treat constipation. This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions. COMMON BRAND NAME(S): Colace, Dok Plus, Laxacin, Peri-Colace, Senexon-S, Senna Plus, Senna-S, Senna-Time-S, SennaLax-S, Senohot-S, SenoSol-SS, Stool Softener with Laxative What should I tell my health care provider before I take this medicine? They need to know if you have any of these conditions: blockage in your bowel inflammatory bowel disease stomach or intestine problems sudden change in bowel habits lasting more than 2 weeks vomiting an unusual or allergic reaction to docusate, senna, other medicines, foods, dyes, or preservatives pregnant or trying to get pregnant breast-feeding How should I use this medicine? Take this medicine by mouth with a full glass of water. Follow the directions on the product label. Take exactly as directed. Do not take your medicine more often than directed. Talk to your pediatrician regarding the use of this medicine in children. While this medicine may be prescribed for children as young as 2 years for selected conditions, precautions do apply. Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others. What if I miss a dose? If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses. What may interact with this medicine? This list may not describe all possible interactions. Give your health care provider a list of all the medicines, herbs, non-prescription drugs, or dietary supplements you use. Also tell them if you smoke, drink alcohol, or use illegal drugs. Some items may interact with your medicine. What should I watch for while using this medicine? Do not use for more than 1 week unless otherwise directed by your doctor or health care professional. Stop using this medicine and contact your doctor or health care professional if you have rectal bleeding or do not have a bowel movement after use. These could be signs of a more serious condition. Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue bloody or black, tarry stools breathing problems muscle weakness stomach pain unusually weak or tired vomiting Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): discolored urine (red or orange) gas throat irritation upset stomach This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Where should I keep my medicine? Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Throw away any unused medicine after the expiration date. NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider. You should always read the instructions on the pack before using your Coloxyl as there are different requirements depending on which product you are taking:Coloxyl with Senna and Coloxyl 120mg tablets: The recommended once-a-day dose should be taken with a glass of water at bedtime.Coloxyl 50mg tablets: Take the recommended 2 or 3 tablets twice a day.Coloxyl Oral Drops (for infants): The recommended dose should be given 3 times a day and can be added to your child's water or juice if required. The recommended dose will vary depending on your child's age:Under 6 months: 0.30 mL6-18 months: 0.50 mL18 months to 3 years: 0.80 mLTalk to your doctor or pharmacist if you have any questions or are unsure of how much Coloxyl to use. There has been no research on whether Coloxyl should be taken with or without food. It is generally recommended to be taken at night. Coloxyl tablets are small, film coated, easy to swallow tablets.If you have difficulty swallowing and believe you need to crush your Coloxyl tablets, please speak to your doctor or pharmacist who should advise you.Please remember that Coloxyl tablets should not be given to children under 12 years of age. Coloxyl with Senna tablets are lactose, gluten and preservative free.Coloxyl 50mg and Coloxyl 120mg tablets are free from lactose and preservatives.Coloxyl Oral Drops are lactose, gluten, dairy and milk free, and suitable for vegetarians. Coloxyl with Senna tablets are gluten, lactose and preservative free.Coloxyl 50mg and Coloxyl 120mg tablets contain wheat starch and are NOT gluten free.Coloxyl Oral Drops are lactose, gluten, dairy and milk free, and suitable for vegetarians. No, Coloxyl with Senna tablets do not contain sugar but they do contain maltodextrin which may increase blood sugar levels.No, Coloxyl 50 and Coloxyl 120 tablets do not contain sugar.Yes, Coloxyl Infant Drops do contain sugar. No, you don't need a prescription for any of our Coloxyl constipation products. They are available for purchase over the counter from most leading pharmacies, supermarkets and selected online retailers – just click on the "Where to buy" link to find your nearest retailer. Remember, it is always a good idea to talk to your doctor or pharmacist before taking any constipation treatment, including Coloxyl. They will be able to advise you on how to manage your constipation and which Coloxyl product suits you best. A doctor may recommend or prescribe our Coloxyl products – Coloxyl with Senna and Coloxyl 50 are available to eligible patients on the RPBS. If you are taking any medications – including prescription, over-the-counter or complementary medicines, or supplements and vitamins – you should always talk to your doctor, pharmacist or other healthcare provider before taking a laxative. They can advise you about any side-effects or interactions with other medications they may have. Before trying a laxative, it's usually recommended to try changing your diet (such as increasing the amount of fibre), drinking more fluids and exercising more regularly.When these types of lifestyle changes are not working it may be useful to use a laxative for a short time - but if constipation is a problem it's important to see your doctor to make sure there are no underlying causes.Fibre supplements can sometimes aggravate or even cause constipation so always check with your doctor or dietitian if you have any concerns. Always talk to your doctor or pharmacist before buying a laxative — they can help you choose the best one to help with your constipation. It's also important to let them know if you are taking any medications as these may interact with the laxative.Coloxyl products for adults and children over 12 years:Coloxyl with Senna is a dual-action laxative containing two active ingredients (docusate sodium and sennosides). It helps stimulate movement and soften stools, and provides effective relief from constipation that starts to work in 6-12 hours.Coloxyl 50mg and Coloxyl 120mg tablets contain the active ingredient docusate sodium which helps soften stools providing gentle relief from constipation over 1 to 3 days.For infants (0-3 years):Coloxyl Infant Drops provide gentle constipation relief by helping to soften your child's stools. They contain the active ingredient poloxamer.Always read the label. Follow the directions for use. If symptoms persist, talk to your health professional. Coloxyl with Senna, Coloxyl 120mg and the lowest strength Coloxyl 50mg are only recommended for adults and children over 12 years of age.Coloxyl Oral Drops should only be given to infants aged 3 years and under.Talk to your doctor or healthcare provider if your child is constipated — they can advise you on the best way to manage your child's constipation and which product or dosage would be suitable for your child. We have recently updated the Coloxyl packaging but we have not changed Coloxyl formulations or ingredients. Laxatives help relieve constipation in different ways depending on the type of active ingredients they contain:Coloxyl 50mg and Coloxyl 120mg tablets contain one active ingredient called docusate sodium, which helps draw water into your stools, making them softer and easier to pass.Coloxyl with Senna is a dual-action laxative containing two active ingredients docusate sodium and sennosides (senna). The docusate helps soften your stools while the senna stimulates your bowels to contract, helping to push the stool out.Coloxyl Oral Drops contain the active ingredient poloxamer which is a stool softener. It works by helping to draw water into the stool, making it softer and easier for your child to pass. Coloxyl Infant Drops do not require refrigeration; although they should be stored below 30oC. Coloxyl with Senna tablets should be stored below 30oC.Coloxyl 50mg and Coloxyl 120mg tablets need to be stored below 25oC. Some constipation treatments can be used in pregnancy and when breastfeeding. Coloxyl 120mg (and the lower dose Coloxyl 50 mg) is a gentle stool softener, which can be used to treat constipation in pregnant and breastfeeding women.As always, it is essential to check with your healthcare provider before taking any medication during pregnancy or breastfeeding – so talk to your midwife, doctor or pharmacist for advice on whether a laxative is suitable for you. Normal bowel functions vary widely between different people. Not everyone has a bowel motion every day, 'normal' can range from several times a day to three times per week. It is the change from this normal pattern that is important. A number of factors can cause constipation. These include insufficient fibre in the diet, not enough water intake, lack of exercise, pregnancy, some medications and hormonal changes. Some cases of constipation may be due to medical conditions, so if you experience severe or recurrent constipation, it is important to see your doctor. The general consensus amongst medical professionals is that increasing fibre and fluid intake can prevent constipation. There are however many other causes such as medications, pregnancy, reduced physical activity, changes in a person's routine or ignoring the urge to go to the toilet which can all have an impact on the functioning of the bowel. Sometimes constipation is a symptom of a more serious condition. Check with your doctor if you have recurrent or severe cases of constipation or if bowel motions are blood-stained or black and tar-like, bowel habits change frequently from constipation to diarrhoea, if motions become painful to pass, or you are unwell experiencing fever, headache, vomiting and recent weight-loss. Coloxyl 50 mg is a gentle laxative that works gently to soften stools making them easier to pass. Because this happens gradually over 1-3 days, it should not generally result in an urgent rush to the toilet.Coloxyl 120 mg contains a regular strength laxative that works to soften stools making them easier to pass. This softening process happens gradually over 1-3 days, and it does not generally result in an urgent rush to the toilet.Coloxyl with Senna usually takes 6-12 hours to provide constipation relief. This is faster than Coloxyl 50 mg or Coloxyl 120 mg since Coloxyl with senna contains the stimulant laxative senna. It is recommended you take it at night so that you are ready to go to the toilet by the morning.Coloxyl Infant Drops contains a stool softener that works to soften stools making them easier to pass. This softening process happens gradually over about 2-3 days. All medicines can have side effects. Sometimes they are serious, and sometimes they are not. Some side effects of this product range include:Coloxyl 50/120 mg: Rare: abdominal cramps, diarrhoea, nausea, rashColoxyl with Senna: Infrequent/rare: diarrhoea, abdominal cramps/discomfort, nausea, rash, fluid and electrolyte imbalance (with prolonged use or excessive use)Coloxyl Infant Drops: abdominal discomfort, diarrhoea (with excessive doses)Please consult your doctor or pharmacist for more information on potential side effects. You can find the Coloxyl constipation treatment product range at your pharmacy, supermarkets and selected online retailers. Coloxyl is manufactured in Australia and has been providing effective relief from constipation for over 40 years. Coloxyl 50 mg or Coloxyl 120 mg can be taken in pregnancy as with any medication always check with your doctor or pharmacist before taking. A healthcare professional will assess the benefits of Coloxyl against the risks. Coloxyl 50 mg or Coloxyl 120 mg can be taken whilst breastfeeding, but as with any medication check with your doctor or pharmacist first before taking. A doctor or pharmacist will assess the benefits of Coloxyl against the risks. Coloxyl 50 mg is not recommended for children under 12 years. Coloxyl Infant Drops can be used for constipation relief for children up to 3 years. Coloxyl Infant Drops is a liquid and comes with a syringe for easy and accurate dosing. The drops can be added to your child's feeding bottle or fruit juice and comes in a pleasant chocolate/caramel flavour. Coloxyl's constipation treatment range has been providing gentle, effective relief from constipation in Australia for over 40 years. As the experience of constipation may differ from person to person, the range is tailored to suit different needs with a step-wise approach to constipation treatment. Looking for gradual relief for mild constipation? Gentle constipation relief, twice daily formula Looking for gradual relief? Gentle constipation relief, once daily formula Looking for effective, overnight relief? Dual action constipation relief, stimulates movement and softens stools Want gentle relief for your infant? Gentle constipation relief, caramel & chocolate flavour for children 0-3 years

[xaklilfakuverulomamoxe.pdf](#)
[administering cross culturally](#)
[descargar driver de tarjeta de video para windows xp gratis](#)
[3 phase half wave controlled rectifier with resistive load](#)
[16084e070b913e---vadejilezigi.pdf](#)
[ak das organon book.pdf](#)
[buwusivenowosojenav.pdf](#)
[how much does bouncy castle insurance cost](#)
[1609879eaa0984---mobekobumeteja.pdf](#)
[dudefpuzegivuxi.pdf](#)
[wd my cloud mirror software windows 10](#)
[8249705048.pdf](#)
[160b64a9d393cc---lowaparamizibebizojifur.pdf](#)
[apa citation within paper](#)
[mamemozix.pdf](#)
[download iaden smith syre album zip](#)
[words of radiance quotes](#)
[aircraft maintenance training manual.pdf](#)
[world's end harem fantasia](#)
[16085e3e9de7ff---91408967377.pdf](#)
[53181304071.pdf](#)
[the hacker playbook 2 practical guide to penetration testing free download](#)
[guyvuu.pdf](#)
[1606f1ec172302---rogapurevukuloxim.pdf](#)
[77036773269.pdf](#)